

## **AMR SEMINAR #63 – SHORT SUMMARY OF CASES:**

- Case 1:** F.51 with constitutional symptoms and 4.3 cm. liver mass.
- Case 2:** M.37 with a firm, solid nodule in the vastus medialis muscle
- Case 3:** F.23 with a 10 cm subcutaneous mass of the anterior chest wall.
- Case 4:** F.39 with a large palpable mass in her right breast.
- Case 5:** F.23 with an endocervical polyp.
- Case 6:** M.21 with enlarged retroperitoneal lymph nodes.
- Case 7:** F.76 with incidental 7.0 cm tumor found in the vicinity of adrenal gland at autopsy.
- Case 8:** F.39 with a palpable nodule located in the soft tissue of the right forearm
- Case 9:** M.59 with a 10 cm. mass in the deep soft tissue of the thigh.
- Case 10:** M.60 with a 3 cm mass on the lesser curvature of the stomach.
- Case 11:** F.43 nursing home resident presented with a fracture of the left proximal tibia.
- Case 12:** F.83 was admitted with a polypoid lesion in the left inguino-femoral fold.
- Case 13:** F.71 presents with a uterine mass 15.3 cm in greatest diameter.
- Case 14:** M.56 with a 1 cm subcutaneous nodule in the thigh.
- Case 15:** F.20 with 3.6 cm mass in left pulmonary hilum.
- Case 16:** F.32 with normal delivery has a 2 cm. necrotic nodule in the placenta.
- Case 17:** F.26 with history of hepatitis B and oral contraceptive use had a liver nodule removed.
- Case 18:** F.32 had a well-circumscribed 3.5 cm. mass removed from her gastric wall.
- Case 19:** M.56 with recurrent pericardial effusion has a biopsy of thick fibrinous material adhering to the pericardium.
- Case 20:** M.76 with 7.0 cm. renal mass.
- Case 21:** M.82 with well-circumscribed 3.1 cm. intrathyroidal nodule.
- Case 22:** F.58 with 2.7 cm. mass of breast.
- Quiz Case #1:** M.43 with enlarged inguinal lymph node.
- Quiz Case #2:** M.68 with 8 cm. soft tissue mass in the lumbar region.
- Quiz Case #3:** M.19 with subpleural ill-defined lung mass extensively infiltrating the overlying pleura.